

Section 1.4

Defining Data Spread

Curriculum Outcomes	Related Activities	Page in Text
<ul style="list-style-type: none"> analyze statistical summaries, draw conclusions, and communicate results about distributions of data 	<ul style="list-style-type: none"> investigate sets of data with a variety of distributions to develop the concept of dispersion, which includes range and variation 	27
<ul style="list-style-type: none"> calculate various statistics using appropriate technology, analyze and interpret the displays, and describe the relationships 	<ul style="list-style-type: none"> investigate sets of data with the same range and mean but different variation, to conclude that variation is independent of range and the location of the data 	30
<ul style="list-style-type: none"> calculate and apply the mean and standard deviation using technology to determine whether a variation makes a difference 	<ul style="list-style-type: none"> compute standard deviations of data sets with a variety of variations and relate standard deviation values to the shape of graphs of those data sets 	29

Section 1.4 -- Pg.27

Dispersion

A measure of the spread of data including the range of the data and the variation within the set of data.

Note: The "range" gives a good indication of the amount of spread, but is a poor indicator of variation within the data. It also provides no information about the location of the central values.

Standard Deviation

A number that describes the spread within a set of data. It represents the average distance a random piece of data is likely to be located from the mean of the data.

-->In other words:

- If there are lots of values far away from the mean, the standard deviation will be greater than if most of the values are close to the mean.

- Note: The standard deviation does not indicate the location of the data but how the data values within a set relate to each other and to the mean.

Refer to Pg.27-29

Example



The manager of a Christmas-tree farm did a study on the time (in hours) needed to sell scotch pines versus white pines once the trees were cut down. The results were as follows;

	TIMES (hrs)								
Scotch Pines	2.15	7.34	9.14	9.50	10.3	10.5	10.7	12.1	21.7
White Pines	2.28	2.46	3.25	4.25	10.2	13.5	14.4	21.2	21.7

Compare the times for the two types of trees by referring to pages 27 - 29:

- 1) Both trees have the same *mean* time of 10.4
- 2) Their *ranges* differ only by 0.2hr,
Scotch Pines Range → 19.6 (2.15 to 21.7)
White Pines Range → 19.4 (2.28 to 21.7)

- 3) Their histograms differ dramatically.
Shade in the appropriate spaces below.



By referring to the *histograms* above, predict which data set would have a **larger** *standard deviation*. →

For a numerical description of the data spread, calculate the Standard Deviation.

First calculate the Standard Deviation for White Pines:

White Pines

Mean \bar{X}	Data Value X	Deviations From the Mean $\bar{X} - X$	Squared Deviations
10.4	2.28	$10.4 - 2.28 = 8.12$	$(8.12)^2 = 65.9$
10.4	2.46	$10.4 - 2.46 = 7.94$	$(7.94)^2 = 63.0$
10.4	3.25	$10.4 - 3.25 = 7.15$	$(7.15)^2 = 51.1$
10.4	4.25	$10.4 - 4.25 = 6.15$	$(6.15)^2 = 37.8$
10.4	10.2	$10.4 - 10.2 = 0.2$	$(0.2)^2 = 0.04$
10.4	13.5	$10.4 - 13.5 = -3.1$	$(-3.1)^2 = 9.61$
10.4	14.4	$10.4 - 14.4 = -4.0$	$(-4.0)^2 = 16.0$
10.4	21.2	$10.4 - 21.2 = -10.8$	$(-10.8)^2 = 116$
10.4	21.7	$10.4 - 21.7 = -11.3$	$(-11.3)^2 = 128$



$\frac{(\text{Sum of Squared Deviations})}{(\text{Number of Data Values})}$ <p>[which is the mean of the squared deviations]</p>	= 54.2
--	--------



The square root of the mean of the squared deviations:	= 7.36
--	--------

Therefore, the Standard Deviation (S_x) for White Pines is $\rightarrow 7.36\text{hr}$

This means that a random piece of data from the White Pines data set is, on average, 7.36hr away from the mean.

Determine the Standard Deviation for Scotch Pines

Review of Steps for calculating Standard Deviation:

Step #1 = Find the \bar{X} of the data

Step #2 = Calculate the distance each data value is from the mean. Square the difference.

Step #3 = Find the \bar{X} of the sum of the squares.

Step #4 = Find the square root of the mean.

Scotch Pines

Mean \bar{X}	Data Value X	Deviations From the Mean $\bar{X} - X$	Squared Deviations $(\bar{X} - X)^2$
10.4		$10.4 - \underline{\hspace{1cm}} =$	$(\underline{\hspace{1cm}})^2 =$
10.4		$10.4 - \underline{\hspace{1cm}} =$	$(\underline{\hspace{1cm}})^2 =$
10.4		$10.4 - \underline{\hspace{1cm}} =$	$(\underline{\hspace{1cm}})^2 =$
10.4		$10.4 - \underline{\hspace{1cm}} =$	$(\underline{\hspace{1cm}})^2 =$
10.4		$10.4 - \underline{\hspace{1cm}} =$	$(\underline{\hspace{1cm}})^2 =$
10.4		$10.4 - \underline{\hspace{1cm}} =$	$(\underline{\hspace{1cm}})^2 =$
10.4		$10.4 - \underline{\hspace{1cm}} =$	$(\underline{\hspace{1cm}})^2 =$
10.4		$10.4 - \underline{\hspace{1cm}} =$	$(\underline{\hspace{1cm}})^2 =$
10.4		$10.4 - \underline{\hspace{1cm}} =$	$(\underline{\hspace{1cm}})^2 =$
10.4		$10.4 - \underline{\hspace{1cm}} =$	$(\underline{\hspace{1cm}})^2 =$



$\frac{(\text{Sum of Squared Deviations})}{(\text{Number of Data Values})}$ <p>[which is the mean of the squared deviations]</p>	$= \underline{23.4}$
--	----------------------



$\sqrt{\frac{(\text{Sum of Squared Deviations})}{(\text{Number of Data Values})}}$	$= \underline{4.84}$
--	----------------------

Therefore, the **Standard Deviation** (S_x) for Scotch Pines is →

This means that a random piece of data taken from the Scotch Pines data set is, on average, _____ hr away from the mean.

Which set of times has more variation, Scotch Pines or White Pines? →

Suppose another type of tree, Douglas fir, produced a set of times with a greater range than white pine. Does that mean this set would automatically have a greater standard deviation? Explain.

Douglas Fir Example: Answer

Even if a data set has a greater range that does not mean that it will have a greater standard deviation.

The two extremes values (the first and last numbers) may be farther away from the mean, however the remaining values might be closer or even at the mean, which would mean that the overall variation might not be greater.

See example: Douglas fir values

0.4, 10.4, 10.4, 10.4, 10.4, 10.4, 10.4, 10.4, 20.4

The range is 20.0 (20.4 - 0.4)

this is greater than for scotch and white pines

The standard deviation is 4.71

this is less than for scotch and white pines

Practice Problem

The amount of snowfall, in centimeters, on eight days were measured and recorded:

6 8 10 15 3 6 4 12

- a) Find the mean snowfall.
- b) Copy and complete the following table for the data:

Mean	Snowfall (Data Value)	Difference from Mean (Mean - Data value)	Squared Deviation (Difference ²)
	3		
	4		
	6		
	6		
	8		
	10		
	12		
	15		

- c) Calculate the mean of the squared difference.
- d) Find the square root of your answer in (c).
What is the standard deviation of the data?

Practice Problem

The amount of snowfall, in centimeters, on eight days were measured and recorded:

6 8 10 15 3 6 4 12

- a) Find the mean snowfall. **8**
- b) Copy and complete the following table for the data:

Mean	Snowfall (Data Value)	Difference from Mean (Mean - Data value)	Squared Deviation (Difference ²)
8	3	8 - 3 = 5	25
8	4	8 - 4 = 4	16
8	6	8 - 6 = 2	4
8	6	8 - 6 = 2	4
8	8	8 - 8 = 0	0
8	10	8 - 10 = -2	4
8	12	8 - 12 = -4	16
8	15	8 - 15 = -7	49

- c) Calculate the mean of the squared difference.

$$118/8 = 14.75$$

- d) Find the square root of your answer in (c).

What is the standard deviation of the data?

$$\sqrt{14.75} = \text{3.84}$$

Classwork/Homework

The hours of TV watched per week by 12 families in Newcastle are:

34 22 28 0 17 30 18 26 28 8 3 24

- a) Find the mean of the data?
- b) Copy and complete the following table for the data:

Mean	TV Watched (Data value)	Difference from Mean (Mean - Data value)	Squared deviation (Difference ²)

- c) Calculate the mean of the squared difference.
- d) Find the square root of your answer in (c).
What is the standard deviation of the data?

Classwork/Homework

The hours of TV watched per week by 12 families in Newcastle are:

34 22 28 0 17 30 18 26 28 8 3 24

- a) Find the mean of the data? 19.8
- b) Copy and complete the following table for the data:

Mean	TV Watched (Data value)	Difference from Mean (Mean - Data value)	Squared deviation (Difference ²)
20	34	20 - 34 = -14	196
20	22	20 - 22 = -2	4
20	28	20 - 28 = -8	64
20	0	20 - 0 = 20	400
20	17	20 - 17 = 3	9
20	30	20 - 30 = -10	100
20	18	20 - 18 = 2	4
20	26	20 - 26 = -6	36
20	28	20 - 28 = -8	64
20	8	20 - 8 = 12	144
20	3	20 - 3 = 17	289
20	24	20 - 24 = -4	16

- c) Calculate the mean of the squared difference.

$$1326/12 = 110.5$$

- d) Find the square root of your answer in (c).

What is the standard deviation of the data?

$$\sqrt{110.5} = 10.53 = 10.5 \text{ or } 11$$

Warm-up

The hours of playing video games per week by 12 random teenagers:

0 7 15 4 6 10 18 25 0 2 0 9

- a) Find the mean of the data?
- b) Copy and complete the following table for the data:

Mean	hrs playing video games (Data value)	Difference from Mean (Mean - Data value)	Squared deviation (Difference ²)

- c) Calculate the mean of the squared difference.
- d) Find the square root of your answer in (c).
What is the standard deviation of the data?

Warm-up



The hours of playing video games per week by 12 random teenagers:

0 7 15 4 6 10 18 25 0 2 0 9

- a) Find the mean of the data? **8**
- b) Copy and complete the following table for the data:

Mean	hrs playing video games (Data value)	Difference from Mean (Mean - Data value)	Squared deviation (Difference ²)
8	0	8 - 0 = 8	64
8	7	8 - 7 = 1	1
8	15	8 - 15 = -7	49
8	4	8 - 4 = 4	16
8	6	8 - 6 = 2	4
8	10	8 - 10 = -2	4
8	18	8 - 18 = -10	100
8	25	8 - 25 = -17	289
8	0	8 - 0 = 8	64
8	2	8 - 2 = 6	36
8	0	8 - 0 = 8	64
8	9	8 - 9 = -1	1

- c) Calculate the mean of the squared difference.

$$692/12 = 57.7$$

- d) Find the square root of your answer in (c).

What is the standard deviation of the data?

$$\sqrt{57.7} = 7.6$$

#4. For each set of data below, create another set of data with the parameters given. Explain what you did to create each set.

(a) masses of rainbow trout caught during a fishing contest:

1.2, 5.1, 1.1, 0.8, 1.3, 1.1, 0.9, 1.3, 1.1

Data in order: 0.8, 0.9, 1.1, 1.1, 1.1, 1.2, 1.3, 1.3, 5.1

Calculate the range and S.D. so that we know what the new set of data should have.

The new set of data should have:

- the same number of values (9)
- a range of 4.3
- a S.D. of the same or lesser value (≤ 1.3)
- a different mean (mean was 1.5)

EXAMPLE:

0.8, 1.5, 1.5, 1.5, 1.5, 1.5, 1.5, 1.5, 5.1

- has 9 numbers
- has a range of 4.3
- has a S.D. of 1.2
- has a mean of 1.8

(b) temperatures recorded at 1-h intervals during the day:

-3, -2, -1, -1, 0, 1, 2, 3, 2, 1, 0, -2

Data in order: -3, -2, -2, -1, -1, 0, 0, 1, 1, 2, 2, 3

Calculate the range and S.D. so that we know what the new set of data should have.

The new set of data should have:

- the same number of values (12)
- a range of 6
- the same mean (mean is 0)
- a standard deviation less than before (< 2)

EXAMPLE:

-3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 3

- the same number of values (12)
- a range of 6
- mean is 0
- the standard deviation is 1

Questions #5 and #6

#5. For each set of data below, create another set of data with the same number of values, the same amount of variation, and the same range, but with a different mean. Explain what you did to create each set.

(a) speeds of cars:

95, 101, 97, 101, 105, 105, 106, 104, 98, 92, 101, 100, 96, 102

-if you change each value by the same amount, the mean will change by that same amount, and the range and S.D. will remain the same.

Example: subtract 5 from each value

90, 96, 92, 96, 100, 100, 101, 99, 93, 87, 96, 95, 91, 97

-the range has not changed (still is 14)

-the S.D. has not changed (still is 4)

-the mean has changed (from 100 to 95)

(b) number of hours Jimmy spent watching TV

1, 3, 0, 1, 9, 8, 3

Example: add one to each data value

2, 4, 1, 2, 10, 9, 4

#6. Find the standard deviation of the set of data shown in each graph.

-To do this: Create a set of data using the histograms

For example

In (a), your data values may be:

5, 5, 5, 6, 6, 6, 7, 7, 7, 8, 8, 8, 9, 9, 9

or

5.5, 5.5, 5.5, 6.5, 6.5, 6.5, 7.5, 7.5, 7.5, 8.5, 8.5, 8.5, 9.5, 9.5, 9.5

-now find the mean, fill out the table, calculate S.D.

Warm-Up

The manager of a Christmas - tree farm did a study on the time (in hours) needed to sell white pines once the trees were cut down. The results are shown below:

2.28 2.46 3.25 4.25 10.2 13.5 14.4 21.2 21.7

- a) Find the mean
- b) Create and fill out the standard deviation table
- c) Find the standard deviation

Warm-Up

6

The manager of a Christmas - tree farm did a study on the time (in hours) needed to sell white pines once the trees were cut down. The results are shown below:

2.28 2.46 3.25 4.25 10.2 13.5 14.4 21.2 21.7

- a) Find the mean 10.36
- b) Create and fill out the standard deviation table
- c) Find the standard deviation 7.4

Classwork/Homework

Do Question #8 Page 31

- Draw and label a box and whisker plot for each
- Draw and label a histogram for each
- Calculate the standard deviation for each

- Compare Group 1 and 2 using all three (box and whisker, histogram, and standard deviations).

Standard Deviation on the TI-83

STEPS:

1. Stat
2. Edit + Enter
3. Input data into a column (preferably L1)
4. Stat + Calc + Enter
5. 2nd function + L (# of column used in 3) + Enter
6. Standard Deviation is σx

EXAMPLE:

The lengths of time, to the nearest minute, required by people in an office pool to type letters are given.

4	7	3	9	8	10	5	10	5	6
8	3	7	1	6	9	8	5	11	8
6	9	4	7	2	8	12	7	9	10

Find the σx of the data.

Classwork

Do Questions Page 32 # 9 - 11

Classwork: Answers

Do Questions Page 32 # 9 - 11

(9) Since each set has the same amount of numbers and their range is the same, their standard deviation will also be the same.

-Adding or subtracting the same amount to each value will cause this situation.

(10) For this question, match the ones that stand out first

d - 0.50 (values are right at the mean)

a - 3.67 (values are spread the furthest from the mean)

c - 1.16 (values are close to the mean)

b - 2.87 (values are further from the mean)

(11) Create 3 data sets that have:

-5 pieces of data

-A mean of 20

-each have a different standard deviation

First, you have to remember how to calculate mean:

-add up the 5 numbers and divide by 5

Second, you need to think of 5 numbers that when added give 100 so that when you divide by 5, you will get 20

Examples:

20,20,20,20,20

50,0,0,0,50

0,10,20,30,40

-each have 5 numbers, each has a mean of 20, and each would have, if calculated, a different S.D.